

TRIUMF Summer Institute 2006  
Collider and Energy Frontier Physics

# Beyond the Standard Model

Lecture 3

Heather Logan  
*Carleton University*

## Plan

Lecture 1      Monday July 17

- Why BSM?
- Supersymmetry

Lecture 2      Monday July 17

- Supersymmetry continued: phenomenology

Lecture 3      Wednesday July 19

- Large extra dimensions: ADD
- Universal extra dimensions; particle spins and UED vs. SUSY

Lecture 4      Thursday July 20

- Deconstruction and the Little Higgs
- T-parity

Lecture 5      Friday July 21

- Warped extra dimensions: RS
- RS and Technicolour

The idea of extra dimensions is not new.

## Kaluza & Klein

In 1919 the mathematician Kaluza was working with Einstein's equations for general relativity.

He discovered that if he put the equations in 5 dimensions, and made the extra dimension a small circle, then what came out in 4 dimensions was ordinary general relativity plus Maxwell's equations (!).

The theoretical physicist Klein (of Klein-Gordon equation fame) is credited with inventing the idea that the extra dimensions could be physically real but curled up very small.

[The tower of excitations of particles in extra dimensions are called Kaluza-Klein (or KK) modes – more in a bit.]

## String theory

String theory is only consistent in 26 dimensions (for bosonic string) or 10 dimensions (for superstring, with fermions and bosons) or maybe 11 dimensions (for supergravity).


The extra dimensions beyond our 3+1 are presumably rolled or folded up very small, so we can't see them.

The “modern age” of extra-dimensional theories started in 1998:

Arkani-Hamed, Dimopoulos & Dvali (ADD) proposed a “solution” of the hierarchy problem: that the scale of quantum gravity is a TeV, not  $10^{18}$  GeV  $\sim M_{\text{Pl}}$  (!)

How could this be possible?

If there are  $\delta$  new extra dimensions with size  $R$ , then below the scale  $R$  the gravitational force law changes from  $1/r^2$  to  $1/r^{2+\delta}$ ! This is just Gauss’s law in more than 3 space dimensions: the gravitational field lines have more room to spread out:



$$V(r) \sim \frac{m_1 m_2}{M_*^{2+\delta}} \frac{1}{r^{1+\delta}} \quad (r \ll R)$$

$$V(r) \sim \frac{m_1 m_2}{M_*^{2+\delta}} \frac{1}{R^\delta r} \equiv \frac{m_1 m_2}{M_{\text{Pl}}^2} \frac{1}{r} \quad (r \gg R)$$

Get a relation between the true higher-dimensional Planck scale  $M_*$  and our four-dim apparent Planck scale  $M_{\text{Pl}} \simeq 10^{18}$  GeV:

$$M_{\text{Pl}}^2 = M_*^{2+\delta} R^\delta$$



But what about Standard Model (non-gravitational) physics?  
We've probed distance scales down to 0.01 fm with colliders  
(1 fm  $\simeq$  size of a proton; corresponds roughly to 1 GeV energy scale)

SM particles can't propagate in the large extra dimensions.  
SM must “live on a brane”: particles confined to a 3+1-dim  
“surface” in the higher-dimensional space, which is no bigger  
than about 1/TeV.

This picture is called **ADD** [after the authors] or **large extra dimensions**.

## ADD phenomenology

The  $\delta$  extra dimensions are compactified – have finite radius  $R$   
A particle propagating in such a space has its momentum in the compactified direction(s) quantized: just like particle-in-a-box.

$$\delta = 1: E^{(n)} = p^{(n)} = 2\pi/\lambda^{(n)} = 2\pi n/L = n/R \quad [\text{definition of } R = L/2\pi]$$

In this case the particles are relativistic, so  $E \propto p$  instead of  $p^2$  – for  $\delta = 1$  the energy levels are evenly spaced  $E \propto n$  (where  $n$  is the mode number).

A 4-dim observer will see the excited states as massive particles, with mass fixed by the quantized momentum in the extra dimension.      Linear spacing of levels,  $M^{(n)} = n/R$ .

$$\delta = 2: E^{(n_1, n_2)} = \sqrt{(p^{(n_1)})^2 + (p^{(n_2)})^2} = \sqrt{(n_1/R_1)^2 + (n_2/R_2)^2}$$

For common  $R_1 = R_2 = R$ , spacing [degeneracy] is:

$$0 [1], 1/R [2], \sqrt{2}/R [1], 2/R [2], \text{ etc.}$$

Energy spacing is  $1/R \sim 1 \text{ meV} - 100 \text{ MeV}$  (for  $\delta = 2$  to 6)

Very dense spacing: Huge multiplicity; can't resolve modes.


Each KK graviton couples with  $1/M_{\text{Pl}}$  strength; sum over huge number of modes makes cross sections respectable at energies near  $M_*$ .

## Two types of collider signal processes:

1. Produce a KK graviton, which escapes from our brane  
 Monojet + missing energy signature


$$pp \rightarrow g G^{(n)}, q G^{(n)}$$

E.g.:



Because we don't have a quantum theory of gravity, we don't really know how to calculate near the cutoff  $M_*$  of the theory.

Leads to some uncertainty in prediction for high-energy cross section:



from Vacavant & Hinchliffe,  
 J. Phys. G **27**, 1839 (2001)




$$pp \rightarrow g G^{(n)}, q G^{(n)}$$

Monojet + missing energy signature

Signal feature: tail at high  $p_T^{\text{miss}}$

plot for LHC from Vacavant & Hinchliffe,  
J. Phys. G **27**, 1839 (2001)



Limits from CDF Run II  
monojet search ( $1.1 \text{ fb}^{-1}$ )


Data agrees with SM

$$\delta = 2 \rightarrow M_D > 1.33 \text{ TeV}$$

...

$$\delta = 6 \rightarrow M_D > 0.88 \text{ TeV}$$


diagram from Beauchemin,  
talk at SUSY'06 conference




Two types of collider signal processes:

2. Produce a pair of SM particles via exchange of a KK graviton

$$pp \rightarrow G^{(n)} \rightarrow f\bar{f}, gg \quad \text{E.g.}:$$



Broad excess in high-invariant-mass Drell-Yan cross section



Many many gravitons – no resonance.  
 Plot is CDF search in  $e^+e^-$  invariant mass:

Data agrees with SM.

Limits from  $D\emptyset$ :

$$\delta = 2 \rightarrow M_D > 1.67 \text{ TeV}$$

...

$$\delta = 6 \rightarrow M_D > 1.14 \text{ TeV}$$

LHC should be sensitive up to  $M_D \simeq 10 \text{ TeV}$ .

**Constraints on ADD:** Astrophysics provides some stringent constraints on the fundamental scale  $M_*$  as a function of the number of extra dimensions  $\delta$ .

	$\delta$			
	2	3	4	5
Supernova Cooling (68)	30	2.5		
Cosmic Diffuse $\gamma$ -Rays:				
Cosmic SNe (69)	80	7		
$\nu\bar{\nu}$ Annihilation (71)	110	5		
Re-heating (72)	170	20	5	1.5
Neutron Star Halo (73)	450	30		
Overclosure of Universe (71)	$6.5/\sqrt{h}$			
Matter Dominated Early Universe (75)	85	7	1.5	
Neutron Star Heat Excess (73)	1700	60	4	1

Astro numbers are rough: depend on scheme for calculating near cutoff  $M_*$ !

**Compare LHC and ILC reach in  $M_*$  [TeV]**

From monojet/mono-photon:

$e^+e^- \rightarrow \gamma + G_n$		2	4	6
LC	$P_{-,+} = 0$	5.9	3.5	2.5
LC	$P_- = 0.8$	8.3	4.4	2.9
LC	$P_- = 0.8, P_+ = 0.6$	10.4	5.1	3.3
$pp \rightarrow g + G_n$		2	3	4
LHC		4 – 8.9	4.5 – 6.8	5.0 – 5.8

From s-channel  $G^{(n)}$ :

		$\sqrt{s}$ (TeV)	$M_H$ (TeV)
LC	$e^+e^- \rightarrow f\bar{f}$	0.5	4.1
	$e^+e^- \rightarrow f\bar{f}$	1.0	7.2
	$\gamma\gamma \rightarrow \gamma\gamma$	1.0	3.5
	$\gamma\gamma \rightarrow WW$	1.0	13.0
	$e\gamma \rightarrow e\gamma$	1.0	8.0
LHC	$pp \rightarrow l^+l^-$	14.0	7.5
	$pp \rightarrow \gamma\gamma$	14.0	7.1


3 tables from Hewett & Spiropulu, hep-ph/0205106

## Black holes at colliders (!)

If the fundamental scale  $M_*$  is as low as a TeV, then LHC collisions will have “trans-Planckian” energies! Offers the possibility to probe quantum gravity directly.

Two colliding partons will come within their mutual higher-dimensional Schwarzschild radius – can form a mini black hole!

Spectrum of black holes produced at LHC with decay to final states tagged with  $e$  or  $\gamma$ :



from Dimopoulos & Landsberg, hep-ph/0106295, via a talk by Landsberg

Is ADD really a solution of the hierarchy problem?

Only “sort of” .

ADD solves the Higgs mass divergence problem.

$$\Delta m^2 \sim (g^2/16\pi^2)\Lambda^2 \text{ is not fine-tuned if } \Lambda \sim \text{TeV}.$$

But, ADD trades a large energy hierarchy for a large size hierarchy: the ratio of  $1/R$  to the fundamental scale  $M_*$ .

$$\delta = 2: R \sim 0.1 \text{ mm} \rightarrow 1/R \sim \text{meV} = 10^{-15} \text{ TeV} \text{ awful!}$$

...

$$\delta = 6: R \sim \text{fm} \rightarrow 1/R \sim 100 \text{ MeV} = 10^{-4} \text{ TeV} \text{ better; still looks tuned.}$$

ADD is not a terrific solution to the hierarchy problem, but experiment will decide.

ADD puts only gravity in the extra dimensions.

What if the SM particles can also propagate in extra dimensions?

This is called Universal Extra Dimensions (UED)

We've probed SM particle interactions up to the 100's of GeV scale: extra dimension must be roughly  $1/(100\text{'s of GeV})$  size or smaller.

Particle-in-a-box for all SM fields:

Get KK towers of SM particles!

UED is not an attempt to solve the hierarchy problem.

However, it gives some very interesting phenomenology and leads the way to other approaches to the hierarchy.

It also provides an interesting “straw-man” model to compare to SUSY.

**Universal extra dimensions:** 5th dimension is a line segment.  
Again get particle-in-a-box KK excitations:  $M^{(n)} = n/R$  (5-dim)

**Gauge boson KK modes:** search for resonances, same as  $Z'$  search: e.g., look in dileptons.

$Z_{KK}, W_{KK}, g_{KK}$  resonances at LHC energies if  $R \sim 1/\text{TeV}$ .  
Tevatron limits in the several-hundred-GeV range from  $ee, \mu\mu$  resonance search.

Indirect constraints from precision EW meas:  $1/R \gtrsim 6 \text{ TeV}$   
Strong limits due to tree-level exchange of KK gauge bosons.  
E.g., new contribution to muon decay with a  $W^{(1)}$  instead of a  $W$  exchange.

**Fermion KK modes:** letting fermions into the 5th dimension complicates things.

A chiral 5-dim fermion corresponds to a left- and right-handed 4-dim fermion!

Need to get rid of the extra components of the zero-modes, so SM fermions stay chiral.

Fermions are dealt with by “**orbifolding**”:

Compactify the 5th dimension on a circle (of radius  $R$ ):  $S_1$

Identify the top half of the circle with the bottom half:  $S_1/Z_2$   
[this is the orbifolding]

Fields odd under the  $Z_2$  orbifold do not have zero modes:

Set up the boundary conditions so that only desired chiral fermions  
(and gauge/Higgs bosons) have zero modes.

Fermion KK modes are vectorlike (Dirac):  $e_L^{(1)}$ ,  $e_R^{(1)}$  separate

5th dimension compactified on  $S_1$ : 5th-dim momentum is conserved!  
 $\sum$  KK number must stay the same in any reaction.

After orbifolding, the 5th-dim momentum is no longer conserved.

There are two special points, the “boundaries” or fixed points of the orbifold.

Lagrangian can include interactions localized at the boundaries: lead to mass splittings between KK modes.

Left over is a “**KK parity**”  $= (-1)^n$  ( $n$  is KK number).


UED with KK parity: Electroweak precision constraints much weaker. No tree-level exchange of level-1 KK excitations.

Limits on KK quark masses  $\sim$  few hundred GeV from direct searches.  
[It's beginning to look a lot like SUSY....]



## UED phenomenology

KK mode masses get radiative corrections from loops of SM particles. **Get splitting in spectrum:**



from Cheng, Matchev, & Schmaltz,  
hep-ph/0205314


This spectrum is for a **common boundary mass** [like  $m_0$  in CMSSM]  
Coloured particles get largest radiative corrections: **get shifted upwards.**

**Lightest odd-parity particle (LKP) is stable:** dark matter candidate; missing energy in decay chains.

**LKP is naturally  $\gamma^{(1)}$  for common boundary terms.**

**This is looking a lot like SUSY!**


# Discovery reach at LHC is comparable to SUSY



from Cheng, Matchev, & Schmaltz, hep-ph/0205314

LHC sensitive up to  $1/R \simeq 1.5$  TeV with  $300 \text{ fb}^{-1}$ .

Because of KK parity, get cascade decay chains:



from Cheng, Matchev, & Schmaltz,  
[hep-ph/0205314](https://arxiv.org/abs/hep-ph/0205314)

Spectrum tends to be more degenerate than SUSY, but collider signals are similar. Jets, leptons, missing  $p_T$


Couplings related to corresponding SM couplings, just like SUSY.  
 KK-odd particles must be pair-produced.

Major difference is particle spins!

SUSY: partners have opposite spin.

UED: partners have same spin.

Sure-fire way to tell you've got UED and not SUSY:  
 detect the 2nd KK level!



$n = 2$  level: even under KK parity. Can singly-produce them as resonances.

from Datta, Kong, & Matchev,  
[hep-ph/0509246](https://arxiv.org/abs/hep-ph/0509246)

Nevertheless, UED gives us a wonderful “straw-man” model to compare to SUSY:

Want to measure spins of SUSY particles to tell they are not KK excitations with some weird spectrum where we've missed the 2nd level!

## Measuring spin at an $e^+e^-$ collider:


Use threshold dependence of cross section.

Consider pair production of slepton pairs; compare to pair production of KK lepton pairs.

Scalar pairs:  $e^+e^- \rightarrow \tilde{\ell}^+\tilde{\ell}^-$  (sleptons):  $\sigma \propto \beta^3$

Fermion pairs:  $e^+e^- \rightarrow l_1^+l_1^-$  (KK leptons):  $\sigma \propto \beta(3 - \beta^2)$

where  $\beta = p/E = \sqrt{1 - 4m^2/s}$  is the velocity of the produced particle.



**Fig. 4.7.** Comparison of spin-0 and spin- $\frac{1}{2}$  particle pair production in  $e^+e^-$  collisions, for particles of mass  $m = 15$  GeV.

Can do a threshold scan:

Get the spin from  $\beta$  dependence of threshold.

Also get very precise mass measurement.


## Another way to measure spin:

Look at angular distribution of pair production.

Consider pair production of slepton pairs; compare to pair production of KK lepton pairs.

Scalar pairs:  $e^+e^- \rightarrow \tilde{\ell}^+\tilde{\ell}^-$  (sleptons):  $\frac{d\sigma}{d\cos\theta} \propto 1 - \cos^2\theta$

Fermion pairs:  $e^+e^- \rightarrow \ell_1^+\ell_1^-$  (KK leptons):  $\frac{d\sigma}{d\cos\theta} \propto 1 + \left(\frac{E^2 - M^2}{E^2 + M^2}\right) \cos^2\theta$



Theory plot: in terms of  
CM angle  $\cos\theta^*$

from Barr, hep-ph/0511115

Plot shows SUSY, UED, and pure “phase space”,  $d\sigma/d\cos\theta =$   
constant for comparison.

It gets a little more tricky because SUSY/KK particles decay:

$$\tilde{\ell}^+ \tilde{\ell}^- \rightarrow \ell^+ \tilde{N}_1 \ell^- \tilde{N}_1 \text{ or } \ell_1^+ \ell_1^- \rightarrow \ell^+ \gamma_1 \ell^- \gamma_1$$

At ILC this is not a problem: can still reconstruct  $\tilde{\ell}^+ \tilde{\ell}^-$  directions.





Figure 3.2.3: Angular distribution of smuons (two entries per event) in the reaction  $e_R^- e_L^+ \rightarrow \tilde{\mu}_R \tilde{\mu}_R \rightarrow \mu^- \tilde{\chi}_1^0 \mu^+ \tilde{\chi}_1^0$ . The hatched histogram represents the false solution.

from Tesla TDR, hep-ph/0106315

At LHC it's more difficult: CM frame is boosted longitudinally.



Measure instead the lepton polar angle in the  $\ell^+ \ell^-$  CM frame.

Still has some sensitivity.

from Barr, hep-ph/0511115

Another method for LHC: look at a decay chain.

Example:

$$\tilde{q} \rightarrow q \tilde{N}_2 \rightarrow q l^\pm \tilde{l}^\mp \rightarrow q l^+ l^- \tilde{N}_1 \text{ in SUSY}$$

$$q_1 \rightarrow q Z_1 \rightarrow q l^\pm l_1^\mp \rightarrow q l^+ l^- \gamma_1 \text{ in UED}$$




diagram from Battaglia, Datta, De Roeck, Kong, & Matchev, hep-ph/0507284


Form  $M_{q\ell}$  invariant mass dist'n with first (near) lepton  
 Shape depends on spin of intermediate particle:  
 $\tilde{N}_2$  in SUSY – spin 1/2;  $Z_1$  in UED – spin 1

Problem: hard to distinguish the first (near) lepton from the second (far) lepton. Tends to wash out spin correlations.




Solution: use a charge asymmetry between  $ql^+$  and  $ql^-$   
 $\widetilde{N}_2$  typically mostly  $\widetilde{W}^0$ : couples to LH fermions / RH antifermions.

Helicity conservation leads to different  $M_{lq}$  shape for  $l^+$  vs.  $l^-$ :



for SUSY




from Barr, hep-ph/0405052

Summing over  $\tilde{q} + \tilde{q}^*$  would wash this out again EXCEPT:  
 LHC is a  $pp$  collider: more  $q$  than  $\bar{q}$  in PDFs.

Make a lepton charge asymmetry:

$$A^{+-} = \frac{s^+ - s^-}{s^+ + s^-}, \quad s^\pm = \frac{d\sigma}{d(M_{\ell^\pm q})}$$

Charge asymmetry depends on  $M_{\ell q}$  differently for **SUSY**, for **UED**, and for pure “phase space” (flat distribution):



from Battaglia, Datta, De Roeck, Kong, & Matchev, hep-ph/0507284

This tests the spin of  $\tilde{N}_2$  or  $Z^{(1)}$ .

There are many other spin observables in other decay chains.

**SUSY:**  $\tilde{g} \rightarrow \tilde{b}_1 \rightarrow \tilde{N}_2 \rightarrow \tilde{\ell}_1 \rightarrow \tilde{N}_1$ . Final state is  $b\bar{b}l^+l^-p_T^{\text{miss}}$ .



**UED:**  $g_1 \rightarrow b_{L1} \rightarrow Z_1 \rightarrow \ell_{R1} \rightarrow \gamma_1$ . Final state is  $b\bar{b}l^+l^-p_T^{\text{miss}}$ .

Choose UED spectrum to match SUSY spectrum!

Have only the spins to distinguish them!

Lepton charge asym. vs.  $M_{b\ell}$   
(softer  $b$ ).

Azimuthal angle dist'n be-  
tween the two  $b$  jets.



from Alves, Éboli, & Plehn, hep-ph/0605067

## Summary:

We talked about:

### ADD

Large extra dimensions, only gravity in the bulk.

Quantum gravity at the TeV scale: mini black holes at colliders!?

### Universal extra dimensions (UED)

1/TeV size extra dimension.

All the SM particles in the bulk (“universal”) – signatures similar to SUSY! [Because of KK parity.]

Challenge: determine spins of new particles to distinguish UED from SUSY.